


## CYPROHEPTADINE HYDROCHLORIDE- cyproheptadine hydrochloride tablet AvPAK

### Cyproheptadine Hydrochloride Tablets, USP Rx Only

#### DESCRIPTION

Cyproheptadine HCl USP is an antihistaminic and antiserotonergic agent.

Cyproheptadine hydrochloride USP is a white to slightly yellowish crystalline solid, with a molecular weight of 350.89, which is soluble in water, freely soluble in methanol, sparingly soluble in ethanol, soluble in chloroform, and practically insoluble in ether. It is the sesquihydrate of 4-(5 *H*-dibenzo [*a,d*]cyclohepten-5-ylidene)-1-methylpiperidine hydrochloride. The molecular formula of the anhydrous salt is C<sub>21</sub>H<sub>21</sub>N•HCl and the structural formula of the anhydrous salt is:



C<sub>21</sub>H<sub>21</sub>N•HCl M.W. 350.89

Cyproheptadine hydrochloride USP is available for oral administration in 4 mg tablets. Inactive ingredients include: lactose monohydrate, magnesium stearate, microcrystalline cellulose, and sodium starch glycolate.

#### CLINICAL PHARMACOLOGY

Cyproheptadine is a serotonin and histamine antagonist with anticholinergic and sedative effects. Antiserotonin and antihistamine drugs appear to compete with serotonin and histamine, respectively, for receptor sites.

#### Pharmacokinetics and Metabolism

After a single 4 mg oral dose of <sup>14</sup>C-labelled cyproheptadine HCl in normal subjects, given as tablets, 2 to 20% of the radioactivity was excreted in the stools. Only about 34% of the stool radioactivity was unchanged drug, corresponding to less than 5.7% of the dose. At least 40% of the administered radioactivity was excreted in the urine. No

detectable amounts of unchanged drug were present in the urine of patients on chronic 12 to 20 mg daily doses. The principle metabolite found in human urine has been identified as a quaternary ammonium glucuronide conjugate of ciproheptadine. Elimination is diminished in renal insufficiency.

## **INDICATIONS AND USAGE**

Perennial and seasonal allergic rhinitis

Vasomotor rhinitis

Allergic conjunctivitis due to inhalant allergens and foods

Mild, uncomplicated allergic skin manifestations of urticaria and angioedema.

Amelioration of allergic reactions to blood or plasma

Cold urticaria

Dermatographism

As therapy for anaphylactic reactions *adjunctive* to epinephrine and other standard measures after the acute manifestations have been controlled.

## **CONTRAINDICATIONS**

### **Newborn or Premature Infants**

This drug should **not** be used in newborn or premature infants.

### **Nursing Mothers**

Because of the higher risk of antihistamines for infants generally and for newborns and prematures in particular, antihistamine therapy is contraindicated in nursing mothers.

### **Other Conditions**

Hypersensitivity to ciproheptadine and other drugs of similar chemical structure.

Monoamine oxidase inhibitor therapy (See **DRUG INTERACTIONS.**)

Angle-closure glaucoma

Stenosing peptic ulcer

Symptomatic prostatic hypertrophy

Bladder neck obstruction

Pyloroduodenal obstruction

Elderly, debilitated patients

## **WARNINGS**

### **Pediatric Patients**

*Overdosage* of antihistamines, particularly in infants and young children, may produce hallucinations, central nervous system depression, convulsions, respiratory and cardiac arrest, and death.

Antihistamines may diminish mental alertness; conversely, particularly, in the young child, they may occasionally produce excitation.

### **CNS Depressants**

Antihistamines may have additive effects with alcohol and other CNS depressants, e.g., hypnotics, sedatives, tranquilizers, antianxiety agents.

### **Activities Requiring Mental Alertness**

Patients should be warned about engaging in activities requiring mental alertness and motor coordination, such as driving a car or operating machinery.

Antihistamines are more likely to cause dizziness, sedation, and hypotension in elderly patients (see **PRECAUTIONS, Geriatric Use**).

## **PRECAUTIONS**

### **General**

Cyproheptadine has an atropine-like action and, therefore, should be used with caution in patients with:

History of bronchial asthma

Increased intraocular pressure

Hyperthyroidism

Cardiovascular disease

Hypertension

### **Information for Patients**

Antihistamines may diminish mental alertness; conversely, particularly, in the young child, they may occasionally produce excitation. Patients should be warned about engaging in activities requiring mental alertness and motor coordination, such as driving a car or operating machinery.

### **Drug Interactions**

MAO inhibitors prolong and intensify the anticholinergic effects of antihistamines.

Antihistamines may have additive effects with alcohol and other CNS depressants, e.g., hypnotics, sedatives, tranquilizers, antianxiety agents.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term carcinogenic studies have not been done with cyproheptadine.

Cyproheptadine had no effect on fertility in a two-litter study in rats or a two generation

study in mice at about 10 times the human dose.

Cyproheptadine did not produce chromosome damage in human lymphocytes or fibroblasts *in vitro*; high doses (10-4M) were cytotoxic. Cyproheptadine did not have any mutagenic effect in the Ames microbial mutagen test; concentrations of above 500 mcg/plate inhibited bacterial growth.

## **Pregnancy**

### **Pregnancy Category B**

Reproduction studies have been performed in rabbits, mice, and rats at oral or subcutaneous doses up to 32 times the maximum recommended human oral dose and have revealed no evidence of impaired fertility or harm to the fetus due to cyproheptadine. Cyproheptadine has been shown to be fetotoxic in rats when given by intraperitoneal injection in doses four times the maximum recommended human oral dose. Two studies in pregnant women, however, have not shown that cyproheptadine increases the risk of abnormalities when administered during the first, second and third trimesters of pregnancy. No teratogenic effects were observed in any of the newborns. Nevertheless, because the studies in humans cannot rule out the possibility of harm, cyproheptadine should be used during pregnancy only if clearly needed.

## **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, and because of the potential for serious adverse reactions in nursing infants from cyproheptadine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother (see **CONTRAINDICATIONS**).

## **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of two have not been established (see **CONTRAINDICATIONS, Newborn or Premature Infants**, and **WARNINGS, Pediatric Patients**).

## **Geriatric Use**

Clinical studies of cyproheptadine HCl tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **WARNINGS, Activities Requiring Mental Alertness**).

## **ADVERSE REACTIONS**

Adverse reactions which have been reported with the use of antihistamines are as follows:

## **Central Nervous System**

Sedation and sleepiness (often transient), dizziness, disturbed coordination, confusion, restlessness, excitement, nervousness, tremor, irritability, insomnia, paresthesias, neuritis, convulsions, euphoria, hallucinations, hysteria, faintness.

## **Integumentary**

Allergic manifestation of rash and edema, excessive perspiration, urticaria, photosensitivity.

## **Special Senses**

Acute labyrinthitis, blurred vision, diplopia, vertigo, tinnitus.

## **Cardiovascular**

Hypotension, palpitation, tachycardia, extrasystoles, anaphylactic shock.

## **Hematologic**

Hemolytic anemia, leukopenia, agranulocytosis, thrombocytopenia.

## **Digestive System**

Cholestasis, hepatic failure, hepatitis, hepatic function abnormality, dryness of mouth, epigastric distress, anorexia, nausea, vomiting, diarrhea, constipation, jaundice.

## **Genitourinary**

Urinary frequency, difficult urination, urinary retention, early menses.

## **Respiratory**

Dryness of nose and throat, thickening of bronchial secretions, tightness of chest and wheezing, nasal stuffiness.

## **Miscellaneous**

Fatigue, chills, headache, increased appetite/weight gain.

To report SUSPECTED ADVERSE REACTIONS contact AvKARE at 1-855-361-3993; email [drugsafety@avkare.com](mailto:drugsafety@avkare.com); or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## **OVERDOSAGE**

Antihistamine overdosage reactions may vary from central nervous system depression to stimulation especially in pediatric patients. Also, atropine-like signs and symptoms (dry mouth; fixed, dilated pupils; flushing, etc.) as well as gastrointestinal symptoms may occur.

*If vomiting has not occurred spontaneously*, the patient should be induced to vomit with syrup of ipecac.

*If patient is unable to vomit*, perform gastric lavage followed by activated charcoal.

Isotonic or ½ isotonic saline is the lavage of choice. Precautions against aspiration must be taken especially in infants and children.

When life threatening CNS signs and symptoms are present, intravenous physostigmine salicylate may be considered. Dosage and frequency of administration are dependent on age, clinical response, and recurrence after response. (See package circulars for physostigmine products.)

*Saline cathartics, as milk of magnesia, by osmosis draw water into the bowel and, therefore, are valuable for their action in rapid dilution of bowel content.*

*Stimulants should not be used.*

Vasopressors may be used to treat hypotension.

The oral LD<sub>50</sub> of cyproheptadine is 123 mg/kg, and 295 mg/kg in the mouse and rat, respectively.

## **DOSAGE AND ADMINISTRATION**

DOSAGE SHOULD BE INDIVIDUALIZED ACCORDING TO THE NEEDS AND THE RESPONSE OF THE PATIENT.

Each tablet contains 4 mg of cyproheptadine hydrochloride.

### **Pediatric Patients**

Age 2 to 6 years

The total daily dosage for pediatric patients may be calculated on the basis of body weight or body area using approximately 0.25 mg/kg/day or 8 mg per square meter of body surface (8 mg/m<sup>2</sup>).

The usual dose is 2 mg (½ tablet) two or three times a day, adjusted as necessary to the size and response of the patient. The dose is not to exceed 12 mg a day.

Age 7 to 14 years

The usual dose is 4 mg (1 tablet) two or three times a day adjusted as necessary to the size and response of the patient. The dose is not to exceed 16 mg a day.

### **Adults**

The total daily dose for adults should not exceed 0.5 mg/kg/day. The therapeutic range is 4 to 20 mg a day, with the majority of patients requiring 12 to 16 mg a day. An occasional patient may require as much as 32 mg a day for adequate relief. It is suggested that dosage be initiated with 4 mg (1 tablet) three times a day and adjusted according to the size and response of the patient.

## **HOW SUPPLIED**

Cyproheptadine Hydrochloride Tablets USP, 4 mg are available as white to off-white round convex tablets, engraved IT above bisect and 68 below bisect on one side with other side is plain, containing 4 mg of cyproheptadine HCl packaged as follows:

NDC 50268-189-15 (10 tablets per card, 5 cards per carton).

PHARMACIST: Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

Dispensed in Unit Dose Package. For Institutional Use Only.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

Manufactured for:

AvKARE

Pulaski, TN 38478

Mfg. Rev. 11/19

AV 03/21 (P)

AvPAK

**PRINCIPAL DISPLAY PANEL - 4 mg Tablet Bottle Label**

NDC 50268-189-15

**Cyproheptadine  
Hydrochloride  
Tablets, USP**

**4 mg**

Rx Only

50 Tablets (5 x 10) Unit Dose



NDC 50268-189-15

**Cyproheptadine  
Hydrochloride  
Tablets, USP**

**4 mg**

Rx Only

50 Tablets (5 x 10) Unit Dose



Each tablet contains 4 mg Cyproheptadine HCl, USP.

Usual Dosage: See package insert for complete prescribing information.

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for:  
AvKARE  
Pulaski, TN 38478



Mfg. Rev. 11/19

AV 03/21 (P)

## CYPROHEPTADINE HYDROCHLORIDE

cyproheptadine hydrochloride tablet

### Product Information

Product Type	Human Prescription Drug	Item Code (Source)	NDC:50268-189(NDC:69367-246)

HUMAN PRESCRIPTION  
DRUG

Item Code  
(Source)

NDC:50268-189(NDC:69367-  
246)



<b>Route of Administration</b>		ORAL		
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
CYPROHEPTADINE HYDROCHLORIDE (UNII: NJ82J0F8QC) (CYPROHEPTADINE - UNII:2YHB6175DO)		CYPROHEPTADINE HYDROCHLORIDE	4 mg	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	2 pieces	
<b>Shape</b>	ROUND	<b>Size</b>	7mm	
<b>Flavor</b>		<b>Imprint Code</b>	IT;68	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:50268-189-15	50 in 1 BOX	03/22/2021	
1	NDC:50268-189-11	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA207555	03/22/2021		

**Labeler -**
AvPAK (832926666)