


MECLIZINE HYDROCHLORIDE- meclizine tablet AvPAK

Meclizine Hydrochloride Tablets, USP

Rx Only

DESCRIPTION

Chemically, meclizinehydrochloride, USP is 1-(p-chloro- α -phenylbenzyl)-4-(m-methylbenzyl) piperazine dihydrochloride monohydrate.



Inactive ingredients for the tablets are: colloidal silicon dioxide, lactose monohydrate, magnesium stearate, microcrystalline cellulose, sodium starch glycolate and talc. The 12.5 mg tablets also contain FD&C Blue #1 Aluminum Lake. The 25 mg tablets also contain D&C Yellow #10 Aluminum Lake.

CLINICAL PHARMACOLOGY

Meclizine hydrochloride is an antihistamine that shows marked protective activity against nebulized histamine and lethal doses of intravenously injected histamine in guinea pigs. It has a marked effect in blocking the vasodepressor response to histamine, but only a slight blocking action against acetylcholine. Its activity is relatively weak in inhibiting the spasmogenic action of histamine on isolated guinea pig ileum.

Pharmacokinetics

The available pharmacokinetic information for meclizine following oral administration has been summarized from published literature.

Absorption

Meclizine is absorbed after oral administration with maximum plasma concentrations reaching at a median T_{max} value of 3 hours post-dose (range: 1.5 to 6 hours) for the tablet dosage form.

Distribution

Drug distribution characteristics for meclizine in humans are unknown.

Metabolism

The metabolic fate of meclizine in humans is unknown. In an in vitro metabolic study using human hepatic microsome and recombinant CYP enzyme, CYP2D6 was found to be the dominant enzyme for metabolism of meclizine.

The genetic polymorphism of CYP2D6 that results in extensive-, poor-, intermediate-and ultrarapid metabolizer phenotypes could contribute to large inter-individual variability in meclizine exposure.

Elimination

Meclizine has a plasma elimination half-life of about 5 to 6 hours in humans.

INDICATIONS AND USAGE

Meclizine hydrochloride tablets are indicated for the treatment of vertigo associated with diseases affecting the vestibular system.

CONTRAINDICATIONS

Meclizine hydrochloride is contraindicated in individuals who have shown a previous hypersensitivity to it.

WARNINGS

Since drowsiness may, on occasion, occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking this drug.

Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma, or enlargement of the prostate gland.

PRECAUTIONS

Pediatric Use

Clinical studies establishing safety and effectiveness in children have not been done; therefore, usage is not recommended in children under 12 years of age.

Pregnancy

Pregnancy Category B. Reproduction studies in rats have shown cleft palates at 25 to 50 times the human dose. Epidemiological studies in pregnant women, however, do not indicate that meclizine increases the risk of abnormalities when administered during pregnancy. Despite the animal findings, it would appear that the possibility of fetal harm is remote. Nevertheless, meclizine, or any other medication, should be used during pregnancy only if clearly necessary.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when meclizine is administered to a nursing woman.

Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of meclizine has not been evaluated. As meclizine undergoes metabolism, hepatic impairment may result in increased systemic exposure of the drug. Treatment with meclizine should be administered with caution in patients with hepatic impairment.

Renal Impairment The effect of renal impairment on the pharmacokinetics of meclizine has not been evaluated. Due to a potential for drug/metabolite accumulation, meclizine should be administered with caution in patients with renal impairment and in the elderly as renal function generally declines with age.

Drug Interactions

There may be increased CNS depression when meclizine is administered concurrently with other CNS depressants, including alcohol, tranquilizers and sedatives (see **WARNINGS**). Based on *in vitro* evaluation, meclizine is metabolized by CYP2D6. Therefore there is a possibility for a drug interaction between meclizine and CYP2D6 inhibitors.

ADVERSE REACTIONS

Anaphylactoid reaction, drowsiness, dry mouth, headache, fatigue, vomiting and, on rare occasions, blurred vision have been reported.

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE, Inc. at 1-855-361-3993; email drugsafety@avkare.com; or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

For the treatment of vertigo associated with diseases affecting the vestibular system, the recommended dose is 25 mg to 100 mg daily, in divided dosage, depending upon clinical response.

HOW SUPPLIED

Meclizine Hydrochloride Tablets, USP **12.5 mg**, are supplied as light blue colored, oval shaped tablets with “AN 441” debossed on one side and plain on the other side.

They are available as follows:

10 Tablets per card, 5 cards per carton: NDC 50268-522-15

Meclizine Hydrochloride Tablets, USP **25 mg**, are supplied as light yellow colored, oval shaped tablets with “AN 442” debossed on one side and plain on the other side.

They are available as follows:

10 Tablets per card, 5 cards per carton: NDC 50268-523-15

Meclizine Hydrochloride Tablets, USP **50 mg**, are supplied as white, oval shaped, partially bisected tablets with “AN 444” debossed on one side and plain on the other side.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Dispensed in Unit Dose package. For Institutional Use Only.

Rx only

Keep out of reach of children

Manufactured for:
AvKARE, Inc.
Pulaski, TN 38478

Mfg. Rev. 04-2018-01
AV Rev. 08/18 (P)
AvPAK

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 50268-522-15

Meclizine Hydrochloride Tablets, USP

12.5 mg

Rx Only

50 Tablets (5 X 10) Unit Dose

5026852215

NDC 50268-522-15

Meclizine Hydrochloride Tablets, USP

12.5 mg

Rx Only

50 Tablets (5 X 10) Unit Dose

5026852215

Each tablet contains:

Meclizine HCl.....12.5 mg

Store at 20 ° to 25 °C (68 ° to 77 °F) [See USP Controlled Room Temperature].

Dispense in tight, light resistant containers as defined in the USP.

DOSAGE AND USE: See accompanying prescribing information.

MOTION SICKNESS: 25 mg to 50 mg daily.

PLEASE KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for:

AvKARE, Inc.

Pulaski, TN 38478

AvPAK

A PRODUCT OF AvKARE

Mfg. Rev. 02-2011 AV 07/13 (P)



NDC 50268-523-15

Meclizine Hydrochloride Tablets, USP

25 mg

Rx Only

50 Tablets (5 X 10) Unit Dose

5026852315

NDC 50268-523-15

Meclizine Hydrochloride Tablets, USP

25 mg

Rx Only

50 Tablets (5 X 10) Unit Dose

5026852315

Each tablet contains:

Meclizine HCl.....25 mg

Store at 20 ° to 25 °C (68 ° to 77 °F) [See USP Controlled Room Temperature].

Dispense in tight, light resistant containers as defined in the USP.

DOSAGE AND USE: See accompanying prescribing information.

MOTION SICKNESS: 25 mg to 50 mg daily.

PLEASE KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for:

AvKARE, Inc.

Pulaski, TN 38478

AvPAK

A PRODUCT OF AvKARE

Mfg. Rev. 02-2011 AV 07/13 (P)



MECLIZINE HYDROCHLORIDE

meclizine tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50268-522(NDC:65162-441)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	12.5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

Product Characteristics

Color	blue (Light)	Score	no score
Shape	OVAL	Size	10 mm
Flavor		Imprint Code	AN;441
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50268-522-15	50 in 1 BOX, UNIT-DOSE	08/08/2013	
1	NDC:50268-522-11	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201451	08/08/2013	

MECLIZINE HYDROCHLORIDE			
meclizine hydrochloride tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50268-523(NDC:65162-442)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)		MECLIZINE HYDROCHLORIDE	25 mg
Inactive Ingredients			
Ingredient Name			Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
TALC (UNII: 7SEV7J4R1U)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
Product Characteristics			

Color	yellow (Light)	Score	no score	
Shape	OVAL	Size	13mm	
Flavor		Imprint Code	AN;442	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50268-523-15	50 in 1 BOX, UNIT-DOSE	08/08/2013	
1	NDC:50268-523-11	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA201451		08/08/2013	

Labeler - AvPAK (832926666)

Revised: 8/2018

AvPAK